

Wirral

Commissioning Policy

CRITERIA

18 March 2024

- 1st April 2019 Published
- 4th June 2019 Amended
- 1st April 2023 Updated To remove superseded policies as part of the Wirral and Merseyside Integrated Care Board (ICB) Commissioning Policy Harmonisation Programme
- September 2023 Updated To include hyperlinks to C&M ICB harmonized policies (Phase 1)
- March 2024 Updated To include hyperlinks to C&M ICB harmonized policies (Phase 2)

Ref:	WIRCCG_2024
Version:	5
Purpose	This document is part of a suite of policies that the Integrated Care Board (ICB) uses to drive its commissioning of healthcare. Each policy in that suite is a separate public document in its own right but will be applied with reference to other policies in that suite.
Supersedes: Wirral Commissioning Policy 2019/2020 issued 1st April 2019 – Updated September 2024	
Author (inc Job Title):	Midlands and Lancashire Commissioning support Unit – Policy Development Programme
Ratified by: (Name of responsible Committee)	NHS Wirral Clinical Commissioning Group - Governing Body (June 2019)
Date Ratified/Updated:	18 March 2024
Date Published and where (Intranet or Website):	18 March 2024 (Website)
Review date:	2024-25
Target audience:	All Wirral & Merseyside NHS organisations and staff

Document control:	ocument control:		
Date: Version Number:		Section and Description of Change	
April 2019	1	Policy ratified by NHS Wirral CCG	
June 2019	2	Policy amended by NHS Wirral CCG	
April 2023 3 ICB por Readed documpart of Co. Global Co.		Policy amended to illustrate removal of policy positions as superseded by Wirral and Merseyside ICB policies 01/04/2023 as referenced and as listed in Appendix 5. Readers should note the following standalone NHS Wirral CCG commissioning policies are documented separately, outside of this document, these will be subject to review and update as part of the ICB's policy harmonisation programme of work: • Continuous Glucose Monitoring • Gluten Free • Subfertility	
September 2023	4	Policy document amended to include hyperlinks to ICB policies (Phase 1).	
18 March 2024 5 Policy document amended to include additional hyperlinks to ICB policies (Phase 2).			

Contents

A.	INTRODUCTION	9
В.	CORE CLINICAL ELIGIBILITY	10
C.	REFERRAL & APPROVAL PROCESS	10
D.	EXCEPTIONALITY	12
E.	PSYCHOLOGICAL DISTRESS	12
F.	PERSONAL DATA (INCLUDING PHOTOGRAPHS)	13
G.	MEDICINES MANAGEMENT	13
н.	EVIDENCE	14
I.	POLICIES	15
1.	Complementary Therapies	15
2.	Dermatology Skin Resurfacing Techniques (including laser dermabrasion and chemical peels) Surgical or Laser Therapy Treatments for Minor Benign Skin Lesions e.g. sebaceous cyst Treatments for Skin Pigment Disorders Surgical/Laser Therapy for Viral Warts (excluding Genital Warts) from Intermediate Tier/ Secondary Care Providers Secondary Care treatment for Acne Vulgaris PMLE (Polymorphic Light Eruption) Treatment - Desensitising Light Therapy using UVB (ultra-violet shortwave) or PUVA (Psoralen combined with UVA)	15 16 16 17
3.	Diabetes Continuous Glucose Monitoring (CGM) Systems for Continuous Glucose Monitoring in Type 1 Diabetes Mellitus Monogenic Diabetes Testing Maturity Onset Diabetes of the Young (MODY)	18
4.	ENT	18 19 20 20

	Septorhinoplasty	21
	Ear Wax removal including microsuction (excluding primary care)	21
5.	Equipment	
	Use of Lycra Suits	22
6.	Fertility	22
	Infertility Treatment for Subfertility e.g. medicines, surgical procedures and assisted conception. This also includes reversal of vasectomy or female sterilisation	n.
7.	General Surgery	22
	Haemorrhoidectomy - Rectal Surgery	22
	Surgery for Treatment of Asymptomatic Incisional and Ventral Hernias	22
	Surgical correction of Diastasis of the Recti	
	Surgery for Asymptomatic Gallstones	
	Lithotripsy for Gallstones	
	Rectopexy and STARR (Stapled Transanal Resection of the Rectum)	23
8	Gynaecology	23
٠.	Surgical Procedures – for the Treatment of Heavy Menstrual Bleeding	
	Hysterectomy with or without Oophrectomy	
5. Equ Use 6. Fer Infe 7. Ger Hae Surry Surry Lith Rec 8. Gyr Surry Surry Surry Surry Surry Surry Sec 9. Mer Inpa Trea Nor Priv stre 10. Neu Bob Trop Fun 11. Opl Upp Low	D&C (dilatation and curettage)	
	Hysteroscopy	
	Fibroid Embolisation / uterine artery embolisation	
	Surgical correction of vaginal/ uterovaginal prolapse	
	Secondary Care follow up of mirena coil insertion	
9	Mental Health	25
٥.	Inpatient Care for Treatment of Chronic Fatigue Syndrome (CFS)	
	Treatment of Gender Dysphoria	
	Non-NHS Drug and Alcohol Rehabilitation (non-NHS commissioned services)	
	Private Mental Health (MH) Care - Non-NHS Commissioned Services: including Psychotherapy, adult eating disorders, general in-patient care, post-traumat	
	stress adolescent mental health	25
10	Neurology	26
10.	Bobath Therapy	
	Trophic Electrical Stimulation for Facial/Bells Palsy	
	Functional Electrical Stimulation (FES)	
11.	Ophthalmology	
	Upper Lid Blepharoplasty - Surgery on the Upper Eyelid	
	Lower Lid Blepharoplasty - Surgery on the Lower Eyelid	
	Surgical Treatments for Xanthelasma Palpebrum (fatty deposits on the eyelids)	27

	Surgery or Laser Treatment for Short Sightedness (myopia) or Long Sightedness (hypermetropia)	
	Cataract Surgery Coloured (irlens) Filters for Treatment of Dyslexia	
	Intra Ocular Telescope for Advanced Age- Related Macular Degeneration	
	Surgical Removal of Chalazion or Meibomian Cysts	
	Surgical treatment for Proptosis/ Dysthyroid eye disease	
	Photodynamic Therapy for ARMD	
	Multifocal (non- accommodative) intraocular lenses	
40		
12.	Oral Surgery Surgical Replacement of the Temporo- Mandibular Joint	20
	Temporo-Mandibular Joint Dysfunction Syndrome & Joint Replacement	
13.	Paediatrics	
	Cranial Banding for Positional Plagiocephaly	29
14.	Plastic & Cosmetic Surgery	29
	Reduction Mammoplasty - Female Breast Reduction	29
	Augmentation Mammoplasty - Breast Enlargement	
	Removal and/or Replacement of Silicone Implants - Revision of Breast Augmentation	
	Mastopexy - Breast Lift	
	Surgical Correction of Nipple Inversion	
	Male Breast Reduction Surgery for Gynaecomastia	
	Hair Removal Treatments including Depilation	
	Laser Treatment or Electrolysis – for Hirsutism.	
	Surgical Treatment for Pigeon Chest	
	Surgical Revision of Scars	
	Apronectomy or Abdominoplasty (Tummy Tuck)	
	Other Skin Excisions/ Body Contouring Surgery e.g. Buttock Lift, Thigh Lift, Arm Lift (Brachioplasty)	
	Treatments to Correct Hair Loss for Alopecia	
	Hair Transplantation	
	Treatments to Correct Male Pattern Baldness	
	Labiaplasty, Vaginoplasty and Hymenorrhaphy	
	Liposuction	
	Rhytidectomy - Face or Brow Lift	
	All procedures undertaken on cosmetic grounds	38
15.	Respiratory	3
. • •	Treatments for Snoring Soft Palate Implants and Radiofrequency	
	Ablation of the Soft Palate	

	Sodium Tetradecyl Sulfate (STS) Injection or 'snoreplasty'	38
	Uvulopalatoplasty and Uvulopalatopharyngopl asy	
	Investigations and treatment for Sleep Apnoea	
	Sleep studies/ Hypersomnia	39
16	Trauma & Orthopaedics	30
10.	Low back pain and sciatica in over 16's Diagnostic, Interventions and Treatments for acute and chronic low back pain. Excluding spinal pathology,	
	radiculopathy and children.	39
	Pharmacological Intervention for lower back pain	
	Pharmacological intervention for sciatica (neuropathic pain in adults)	
	Radiofrequency Facet Joint Denervation	
	Fusion	
	Epidural Injection	
	Spinal Decompression	
	Endoscopic Laser Foraminoplasty	
	Peripheral Nerve-field Stimulation (PNFS) for Chronic Low Back Pain	
	Endoscopic Lumbar Decompression	
	Percutaneous Disc Decompression using Coblation for Lower Back Pain	43
	Non-Rigid Stabilisation Techniques	
	Lateral (including extreme, extra and direct lateral) Interbody Fusion in the Lumbar Spine	
	Percutaneous Intradiscal Laser Ablation in the Lumbar Spine	
	Transaxial Interbody Lumbosacral Fusion	
	Therapeutic Endoscopic Division of Epidural Adhesions	
	Automated Percutaneous Mechanical Lumbar Discectomy	
	Prosthetic Intervertebral Disc Replacement in the Lumbar Spine	44
	Bone Morphogenetic Proteins - Dibotermin Alfa; Eptotermin Alpha	44
	Surgery for Trigger Finger	45
	Hyaluronic Acid and Derivatives Injections for Peripheral Joint Pain	
	Secondary Care Administered Steroid Joint Injections	
	Dupuytren's Disease Palmar Fasciectomy/Needle Faciotomy	
	Radiotherapy Collagenase Injections for Dupuytren's Disease	45
	Dupuytren's Disease Surgical treatment	
	Dupuytrens Contracture – conservative treatment	
	Hip and Knee Replacement Surgery & Hip Resurfacing	
	Diagnostic Arthroscopy for Arthritis of the Knee	
	Arthroscopic Lavage and Debridement for Osteoarthritis of the Knee	
	Patient Specific Total Knee Replacement	
	Surgical Treatment for Carpal Tunnel Syndrome	
	Nerve Conduction Studies for Carpal Tunnel Syndrome	
	Surgical Removal of Mucoid Cysts at Distal Inter Phalangeal Joint (DIP)	47

	Surgical Removal of Ganglions	47
	Hip Arthroscopy for Femoro–Acetabular Impingement	47
	Surgical Removal of Bunions/Surgery for Lesser Toe Deformity	48
	Surgical Treatment of Morton's Neuroma	48
	Surgical Treatment of Plantar Fasciitis	48
	Treatment of Tendinopathies Extracorporeal Shock Wave Therapy Autologous Blood or Platelet Injection	49
	Injections for Tendonitis (Jumpers Knee)	
	Shoulder Arthroscopy (including arthroscopic shoulder decompression for subacromial shoulder pain)	49
	Hip Injections	
17.	Urology	50
	Circumcision	
	Penile Implant: A Surgical Procedure to Implant a Device into the Penis	50
	Male sterilisation under General Anaesthetic	
	Reversal of Male Sterilisation	
	ESWT (extracorporeal shockwave therapy) for Prostadynia or Pelvic Floor Syndrome	50
	Hyperthermia Treatment for Prostadynia or Pelvic Floor Syndrome	
	Surgery for Prostatism	
	Surgical treatment for Hydroceles – adults and children	
	Surgical removal of benign epididymal cysts	51
18.	Vascular	52
	Surgery for Extreme Sweating	52
	Hyperhydrosis – all areas	52
	Surgical Resection Endoscopic Thoracic Sympathectomy	52
	Chelation Therapy for Vascular Occlusions	
	Varicose Veins Interventional Treatments e.g. endothermal ablation, foam sclerotherapy and surgery	52
19.	Other	52
	Botulinum Toxin A & B Used in several types of procedures e.g. to treat muscle disorders, excessive sweating hyperhidrosis) and migraine	52
	Correction of privately funded treatment	55
J.	Appendix 1 – Cataract Referral Guide	56
L.	Appendix 2 – IEFR Process	57
M.	Appendix 3 – IFER Panel Contact Details	58
N.	Appendix 4 – Fusion Surgery – Clinical exceptions permitted	59

A. INTRODUCTION

Wirral CCG are legally obliged to have in place and publish arrangements for making decisions and adopting policies on whether particular health care interventions are to be made available in Wirral. This document is intended to be a statement of such arrangements made by the Wirral CCGs and act as a guidance document for patients, clinicians and other referrers in primary and secondary care. It sets out the eligibility criteria under which Wirral CCGs will commission the service, either via existing contracts or on an individual basis. It gives guidance to referrers on the policies of the CCGs in relation to the commissioning of procedures of low clinical priority, thresholds for certain treatment and those procedures requiring individual approval.

In making these arrangements, the Wirral CCGs have had regard to relevant law and guidance, including their duties under the National Health Service Act 2006, the Health and Social Care Act 2012 and the National Health Service Commissioning Board and Clinical Commissioning Groups (Responsibilities and Standing Rules) Regulations 2012; the Joint Strategic Needs Assessment; and relevant guidance issued by NHS England.

The Wirral CCGs have a duty to secure continuous improvement in the quality of services and patient outcomes but are also under a duty to exercise their functions effectively, efficiently and economically. Therefore, health benefits must be maximised from the resources available. As new services become available, demand increases and procedures that give maximum health gain must be prioritised. This means that certain procedures will not be commissioned by CCGs unless exceptional clinical grounds can be demonstrated. The success of the scheme will depend upon commitment by GPs and other clinicians to restrict referrals falling outside this protocol.

The NHS Standard Contract requires that the provider must manage referrals in accordance with the terms of any Prior Approval Scheme. If the provider does not comply with the terms of any Prior Approval Scheme in providing a service, the commissioners will not be liable to pay for that service. This includes compliance with terms SC28 to SC31 of the contract which specifically reference procedures included in this policy.

CCGs will not pay for activity unless it meets the criteria set out in the document or individual approval has been given and the Referral and Approval Process as set out has been followed. This prior approval scheme will be incorporated into all NHS standard NHS contracts agreed by CCGs. Compliance with this policy will be monitored via regular benchmarking reports and case note audits.

To support this approach a set of Core Clinical Eligibility Criteria have been developed and are set out below; patients may be referred in accordance with the referral process if they meet these criteria. In some limited circumstances, a 'Procedure of Lower Clinical Priority' (PLCP) may be the most clinically appropriate intervention for a patient. In these circumstances, agreed eligibility criteria have been established and these are explained in the later sections of the document, if the criteria are met the procedure will be commissioned by the CCG.

B. CORE CLINICAL ELIGIBILITY

Patients may be referred in accordance with the referral process where they meet any of the following Core Clinical Eligibility criteria:

- All NICE Technology Appraisals will be implemented.
- In cancer care (including but not limited to skin, head and neck, breast and sarcoma) any lesion that has features suspicious of malignancy, must be referred to an appropriate specialist for urgent assessment under the 2-week rule.
- Reconstructive surgery post cancer or trauma including burns.
- Congenital deformities: Operations on congenital anomalies of the face and skull are usually available on the NHS. Some conditions are considered highly specialised and are commissioned in the UK through the National Specialised Commissioning Advisory Group (NSCAG). As the incidence of some cranio-facial congenital anomalies is small and the treatment complex, specialised teams, working in designated centres and subject to national audit, should carry out such procedures.
- Tissue degenerative conditions requiring reconstruction and/or restoring function e.g. leg ulcers, dehisced surgical wounds, necrotising fasciitis. Any patient who needs urgent treatment will always be treated.
- No treatment is completely ruled out if an individual patient's circumstances are exceptional. Requests for consideration of exceptional circumstances should be made to the patient's responsible CCG see the exceptionality criteria in this policy and the contact details at Appendix 1.
- Children under 16 years are eligible for surgery to alter appearance, improve scars, excise facial or other body lesions, where such conditions cause obvious psychological distress.

C. REFERRAL & APPROVAL PROCESS

Interventions specified in this document are not commissioned unless clinical criteria are met, except in exceptional circumstances. Where clinical criteria are met treatment identified will form part of the normal contract activity.

If a General Practitioner/Optometrist/Dentist considers a patient might reasonably fulfil the eligibility criteria for a Procedure of Lower Clinical Priority, as detailed in this document (i.e. they meet the specific criteria listed for each treatment) the General Practitioner/Optometrist/Dentist should follow the process for referral. If in doubt over the local process, the referring clinician should contact the General Practitioner. Failure to comply with the local process may delay a decision being made. The referral letter should include specific information regarding the patient's potential eligibility.

Diagnostic procedures to be performed with the sole purpose of determining whether or not a Procedure of Lower Clinical Priority is feasible should not be carried out unless the eligibility criteria are met or approval has been given by the CCG or GP (as set out in the approval process of the patients responsible CCG) or as agreed by the CCG as an exceptional case.

The referral process to secondary care will be determined by the responsible CCGs. Referrals will either:

Have received prior approval by the CCG.

OR

Clearly state how the patient meets the criteria.

OR

Be for a clinical opinion to obtain further information to assess the patient's eligibility.

GPs should not refer unless the patient clearly meets the criteria as this can raise unrealistic expectations for the patient and lead to disappointment. In cases where there may be an element of doubt the GP should discuss the case with the IFR Team in the first instance.

If the referral letter does not clearly outline how the patient meets the criteria, then the letter should be returned to the referrer for more information and the CCG notified. Where a GP requests only an opinion the patient should not be placed on a waiting list or treated, but the opinion given to the GP and the patient returned to the GP's care, in order for the GP to make a decision on future treatment.

The secondary care consultant will also determine whether the procedure is clinically appropriate for a patient and whether the eligibility criteria for the procedure are fulfilled or not and may request additional information before seeing the patient. Patients who fulfil the criteria may then be placed on a waiting list according to their clinical need. The patient's notes should clearly reflect exactly how the criteria were fulfilled, to allow for case note audit to support contract management. Should the patient not meet the eligibility criteria this should be recorded in the patient's notes and the consultant should return the referral back to the GP with a copy to the CCG, explaining why the patient is not eligible for treatment.

Should a patient not fulfil the clinical criteria but the referring clinician is willing to support the application as <u>clinically exceptional</u>, the case can be referred to the IFR Team for assessment contact details for the IFR team can be found in Appendix 1.

Where the treatment has changed in the middle of a care pathway and a decision to treat has been made based on the old criteria the treatment can be completed i.e. if the patient has been listed for surgery. Where a clinical decision as to the nature of treatment has not yet been made then the new criteria should be applied with immediate effect.

D. EXCEPTIONALITY

In dealing with exceptional case requests for an intervention that is considered to be a poor use of NHS resources, the Wirral CCGs have endorsed through the CCG Alliance the following description of exceptionality contained in a paper by the NW Medicines and Treatment Group:

• The patient has a clinical picture that is significantly different to the general population of patients with that condition **and as a result of that difference**; the patient is likely to derive greater benefit from the intervention than might normally be expected for patients with that condition.

The Wirral CCGs are of the opinion that exceptionality should be defined solely in clinical terms. To consider social and other non-clinical factors automatically introduces inequality, implying that some patients have a higher intrinsic social worth than others with the same condition. It runs contrary to a basic tenet of the NHS namely, that people with equal need should be treated equally. Therefore non-clinical factors will not be considered except where this policy explicitly provides otherwise.

In essence, exceptionality is a question of equity. The CCG must justify the grounds upon which it is choosing to fund treatment for a particular patient when the treatment is unavailable to others with the condition.

E. PSYCHOLOGICAL DISTRESS

Psychological distress alone will not be accepted as a reason to fund surgery except where this policy explicitly provides otherwise. Psychological assessment and intervention may be appropriate for patients with severe psychological distress in respect of their body image but it should not be regarded as a route into aesthetic surgery.

Unless specifically stated otherwise in the policy, any application citing psychological distress will need to be considered as an IFR. Only very rarely is surgical intervention likely to be the most appropriate and effective means of alleviating disproportionate psychological distress. In these cases ideally an NHS psychologist with expertise in body image or an NHS Mental Health Professional (depending on locally available services) should detail all treatment(s) previously used to alleviate/improve the patient's psychological wellbeing, their duration and impact. The clinician should also provide evidence to assure the IFR Panel that a patient who has focused their psychological distress on some particular aspect of their appearance is at minimal risk of having their coping mechanism removed by inappropriate surgical intervention.

F. PERSONAL DATA (INCLUDING PHOTOGRAPHS)

In making referrals to the IFR Team, clinicians and other referrers in primary and secondary care should bear in mind their obligations under the Data Protection Act 1998 and their duty of confidence to patients. Where information about patients (including photographs) is sent to the IFR Team and is lost or inadvertently disclosed to a third party before it is safely received by the IFR Team, the referrer will be legally responsible for any breach of the Data Protection Act 1998 or the law of confidence.

Therefore, please consider taking the following precautions when using the Royal Mail to forward any information about patients including photographic evidence: Clearly label the envelope to a named individual i.e. first name & surname, and job title.

Where your contact details are not on the items sent, include a compliment slip indicating the sender and their contact details in the event of damage to the envelope or package.

Use the Royal Mail Signed for 1st Class service, rather than the ordinary mail, to reduce the risk of the post going to the wrong place or getting lost.

Information in Payment: Costs incurred for photographic evidence will be the responsibility of the referrer. Photographic evidence is often required in cases which are being considered on exceptionality. They are reviewed by clinical member/s of the IFR team only.

G. MEDICINES MANAGEMENT

Prior approval for treatment should always be sought from the responsible Medicine Management Team when using medicines as follows:

- Any new PbR excluded drug where the drug has not yet been approved/prioritised for use in agreement with the local CCG.
- Any existing PbR excluded drugs to be used outside of previously agreed clinical pathways/indication.
- Any PbR excluded drugs that are being used out with the parameters set by NICE both in terms of disease scores or drug use. It must not be assumed that a new drug in the same class as one already approved by NICE can be used, this must be subject to the process in Point 1.
- Any drug used out with NICE Guidance (where guidance is in existence).
- Any proposed new drug/new use of an existing drug (whether covered by NICE or PBR excluded or not) should first be approved by the relevant Area Medicines Management Committee, and funding (where needed) agreed in advance of its use by the relevant CCG.
- Any medicines that are classed by the CCG as being of limited clinical value.
- · Any medicines that will be supplied via a homecare company agreement.

The Clinical Commissioning Group does not expect to provide funding for patients to continue treatment commenced as part of a clinical trial. This is in line with the Medicines for Human Use (Clinical Trials) Regulations 2004 and the Declaration of Helsinki which stipulates that the responsibility for ensuring a clear exit strategy from a trial, and that those benefiting from treatment will have ongoing access to it, lies with those conducting the trial. This responsibility lies with the trial initiators indefinitely.

NOTE: Funding for all solid and haematological cancers are now the responsibility of NHS England.

Conditions & Interventions: The conditions & interventions have been broken down into speciality groups.

GPs should only refer if the patient meets the criteria set out or individual approval has been given by the CCG as set out in the CCG's process as explained above. Requests for purely cosmetic surgery will not be considered except where this policy explicitly provides otherwise. Patients meeting the core clinical eligibility criteria set out above can be referred, all other referrals should be made in accordance with the specified criteria and referral process. The CCG may request photographic evidence to support a request for treatment.

From time to time, CCGs may need to make commissioning decisions that may suspend some treatments/criteria currently specified within this policy.

Where CCGs have variations in their local clinical policies/pathways or clinical thresholds then this will be highlighted in the comments section indicating there is a local CCG addendum.

H. EVIDENCE

At the time of publication the evidence presented was the most current available. Where reference is made to publications over five years old, this still represents the most up to date view.

I. POLICIES

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
1.	Complementary Th	erapies		
1.1	Complementary Therapies	This policy has been superseded by NHS Wir CMICB Clin067 – Complementary and alternat		
2.	Dermatology			
2.1	Skin Resurfacing Techniques (including laser dermabrasion and chemical peels)	Only be commissioned in the following circumstances: Severe scarring following: Acne once the active disease is controlled. Chicken pox. OR Trauma (including post-surgical).	Modernisation Agency's Action on Plastic Surgery 2005. Hædersdal, M., Togsverd-Bo, K., & Wulf, H. (2008). Evidence-based review of lasers, light sources and photodynamic therapy in the treatment of acne vulgaris. Journal of the European Academy of Dermatology and Venereology, 22, 267–78.	
		Procedures will only be performed on the head and neck area. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core" surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG.	Department of Dermatology, Bispebjerg Hospital, University of Copenhagen, Copenhagen, Denmark. Collated on NHS evidence website suggests that short-term efficacy from optical treatments for acne vulgaris with the most consistent outcomes for PDT. www.evidence.nhs.uk Interim Gender Dysphoria Protocol & Service Guidelines 2013/14.	

WIRCCG_2024 – Commissioning Policy Version 5, 18 March 2024

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
			NHS England interim protocol NHS England (2013)	
			Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	
2.2	Surgical or Laser Therapy Treatments for Minor Benign Skin Lesions e.g. sebaceous cyst	This policy has been superseded by NHS Wir CMICB Clin005 – Benign skin lesions	ral & Merseyside ICB Policy:	
2.4	Treatments for Skin Pigment Disorders	This policy has been superseded by NHS Wir CMICB Clin009 – Camouflage Treatment for S		
2.5	Surgical/Laser Therapy for Viral Warts (excluding Genital Warts) from Intermediate Tier/	Will be commissioned in any of the following circumstances: • Severe pain substantially interfering with functional abilities.	Modernisation Agency's Action on Plastic Surgery 2005. Nongenital warts: recommended approaches to management Prescriber	Most viral warts will clear spontaneously or following application of topical treatments.
	Secondary Care Providers	Persistent and spreading after 2 years and refractive to at least 3 months of primary care or community treatment.	2007 18(4) p33-44. Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery.	65% are likely to disappear spontaneously within 2 years.
		 Extensive warts (particularly in the immune-suppressed patient). Facial warts. 	Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service	There are numerous OTC preparations available.
		Patients with the above exceptional symptoms may need specialist assessment, usually by a dermatologist.	patient.co.uk/doctor/viral-warts-excluding-verrucae http://www.patient.co.uk/doctor/verrucae	Community treatments such a cryosurgery, curettage, prescription only topical treatment should be considered before referral

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
				to secondary care.
2.6	Secondary Care treatment for Acne Vulgaris	 Will be commissioned in any of the following circumstances: Patient has severe acne that is unresponsive to oral antibacterials Patient has moderate to severe acne that is partially unresponsive to treatment that is starting to scar Patients with acne who have failed two full courses of oral antibiotic treatment combined with appropriate topical treatment for a minimum of 6 months Patients with severe nodulo-cystic, conglobate acne Patients at risk of post-inflammatory hyperpigmentation Patients with associated and severe psychological symptoms regardless of severity of acne Patients that do not meet this criteria should be managed in Primary Care. 	https://cks.nice.org.uk/acne-vulgaris http://www.nhs.uk/conditions/acne/pages/t reatmentoptions.aspx	ACNE VULGARIS.docx
2.7	PMLE (Polymorphic Light Eruption) Treatment - Desensitising Light Therapy using UVB (ultra-violet shortwave) or PUVA (Psoralen combined with UVA)	Will be commissioned if ALL of the following criteria are met: • Diagnosis by Dermatology Consultant • Severe with symptoms causing significant functional impairment (Symptoms preventing the patient fulfilling vital work or educational responsibilities - Symptoms	http://www.bad.org.uk/shared/get-file.ashx?id=117&itemtype=document http://www.nhs.uk/conditions/polymorphic-light-eruption/Pages/Introduction.aspx	Clinical discussion with the patient should include educating patients not to use sunbeds as an alternative. It is not comparable to desensitising light therapy and carries additional health risks.

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		preventing the patient carrying out vital domestic or carer activities) • Symptoms remain severe despite preventative treatments • Light therapy deemed likely to make significant improvement to patients symptoms		
3. C	Diabetes			
3.1	Continuous Glucose Monitoring (CGM) Systems for Continuous Glucose Monitoring in Type 1 Diabetes Mellitus	This policy has been superseded as follows: At the NHS Wirral & Merseyside ICB Board M commissioning polices in respect of CGMs be and NG28 be adopted.		
3.2	Monogenic Diabetes Testing Maturity Onset Diabetes of the Young (MODY)	This policy has been superseded by NHS Wir CMICB Clin031 – Monogenic Diabetes Testing	•	
4. E	NT			
4.1	Adenoidectomy	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:	
		CMICB Clin002 – Adenoidectomy		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
4.2	Pinnaplasty – for Correction of Prominent Ears	May be commissioned in the following circumstances: Surgical "correction" of prominent ear(s) only when all of the following criteria are met: 1. Referral only for children aged up to 18 years at the time of referral. AND 2. With very significant ear deformity or asymmetry. AND 3. Patients present with significant detrimental impact on child's ability to lead a normal life Patients not meeting these criteria should not be routinely referred for surgery. Incisionless otoplasty is not commissioned.	Pinnaplasty Department of Health (2007). Local PCT consensus - review conducted 2007. Modernisation Agency's Action on Plastic Surgery 2005. IPG 422: Incisionless otoplasty NICE 2012. http://www.rcseng.ac.uk/healthcare-bodies/docs/published-guides/pinnaplasty Royal College of Surgeons (2013).	
4.3	Insertion of Grommets for Glue Ear (otitis media with effusion)	This policy has been superseded by NHS Will CMICB Clin023 – Grommets for glue ear in chi		,

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
4.4	Tonsillectomy for Recurrent Tonsillitis (excluding peritonsillar abscess) Adults and Children	This policy has been superseded by NHS Wir CMICB Clin046 – Tonsillectomy	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin046 – Tonsillectomy	
4.5	Surgical Remodelling of External Ear Lobe	This policy has been superseded by NHS Wir CMICB Clin45 – Split (cleft) Earlobe, surgical re		
4.6	Use of Sinus X-ray	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin44 – Sinus X-Ray		
4.7	Rhinoplasty - Surgery to Reshape the Nose	 This procedure is NOT available under the NHS on cosmetic grounds. Only commissioned in any of the following circumstances: Objective nasal deformity caused by trauma. Problems caused by obstruction of nasal airway. Correction of complex congenital conditions e.g. cleft lip and palate. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core" surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG. 	Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013) Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	Patients with isolated airway problems (in the absence of visible nasal deformity) may be referred initially to an Ear Nose and Throat (ENT) consultant for assessment and treatment.

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
4.8	Surgery of Laser Treatment of Rhinophyma	This policy has been superseded by NHS Wir CMICB Clin41 - Rhinophyma, surgical manage	•	
4.9	Septorhinoplasty	Only commissioned where: patient has a deviated septum causing significant and persistent nasal blockage AND septoplasty alone will not improve functional impairment OR significant symptoms post trauma/cancer treatment/ severe congenital abnormality This procedure is not commissioned for cosmetic reasons	http://www.lnwh.nhs.uk/services/a-z-services/e/ent-ear-nose-and-throat/ent-operations/nose-operations/septorhinoplasty/	
4.10	Ear Wax removal including microsuction (excluding primary care)	 Only commissioned where: Perforated ear drum OR Otitis Externa OR Hearing loss and all other methods of wax removal have been tried and failed OR Enable inspection of ear drum due to clinical concern of other pathologies and other methods of wax removal have failed OR Clinical risk of other methods of removal 	http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2907972/	Ear wax removal should be managed in primary care and does not require onward referral.

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
5.	Equipment			
5.1	Use of Lycra Suits	This policy has been superseded by NHS Will CMICB Clin071 – Lycra™ Suits and Orthotics		
6.	Fertility			
6.1	Infertility Treatment for Subfertility e.g. medicines, surgical procedures and assisted conception. This also includes reversal of vasectomy or female sterilisation.	See Wirral Subfertility Policy.	CG156 Fertility: Assessment and treatment for people with fertility problems – NICE 2013. Contraception – sterilization – NICE Clinical Knowledge Summaries 2012 http://cks.nice.org.uk/contraception-sterilization#!scenario	Individual CCG addendums apply.
	General Surgery			
7.1	Haemorrhoidectomy - Rectal Surgery Removal of Haemorrhoidal Skin Tags	This policy has been superseded by NHS Will CMICB Clin024 – Haemorrhoids, surgical ma		
7.2	Surgery for Treatment of Asymptomatic Incisional and Ventral Hernias	This policy has been superseded by NHS Will CMICB Clin083 – Minimally symptomatic ing		
	Surgical correction of Diastasis of the Recti	This policy has been superseded by NHS Will CMICB Clin014 – Diastasis (divarication) of the control of the con		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments	
7.3	Surgery for Asymptomatic Gallstones		This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin021 – Gallstones (Asymptomatic), Surgical Management		
7.4	Lithotripsy for Gallstones	Lithotripsy not routinely commissioned.		Lithotripsy rarely performed as rate of recurrence is high.	
7.5	Rectopexy and STARR (Stapled Transanal Resection of the Rectum)	This policy has been superseded by NHS Wir CMICB Clin108 – Rectal Prolapse (Internal or E			
8. 0	Synaecology				
8.1	Surgical Procedures – for the Treatment of Heavy Menstrual Bleeding Hysterectomy with or without Oophrectomy	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB_Clin026 - Heavy Menstrual Bleeding, Hysterectomy			
8.2	D&C (dilatation and curettage)	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin025 - Heavy Menstrual Bleeding, Dilatation and Curettage			
8.3	Hysteroscopy	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin076 – Heavy menstrual bleeding – hysteroscopy policy			
8.4	Fibroid Embolisation / uterine artery embolisation	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin075 – Fibroids (myoma, leiomyoma), uterine artery embolisation (UAE)			
8.5	Surgical correction of vaginal/	This will only be commissioned if: symptomatic prolapse (including those	http://www.nhs.uk/conditions/Prolapse-of-the-uterus/Pages/Introduction.aspx		

WIRCCG_2024 – Commissioning Policy Version 5, 18 March 2024

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	uterovaginal prolapse	combined with urethral sphincter incompetence or faecal incontinence) or Prolapse combined with urethral sphincter incompetence/ urinary incontinence or faecal incontinence AND Evidence that conservative management has been trialed in primary care and failed including; lifestyle modification, pelvic floor exercises, vaginal pessary, local oestrogen creams and oral treatments unless contraindicated, AND Symptoms cause significant functional impairment (Symptoms preventing the patient fulfilling vital work or educational responsibilities - Symptoms preventing the patient carrying out vital domestic or carer activities) AND Patient has confirmed that they do want surgery to correct		
8.6	Secondary Care follow up of mirena coil insertion	This policy has been superseded by NHS Wir CMICB Clin078 – Intrauterine devices: sec		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments		
9. N	. Mental Health					
9.1	Inpatient Care for Treatment of Chronic Fatigue Syndrome (CFS)	CMICB Clin066 – Chronic fatigue syndrome	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin066 – Chronic fatigue syndrome/Myalgic Encephalomyelitis (CFS/ME): Inpatient Management			
9.2	Treatment of Gender Dysphoria	This policy has been superseded by NHS Wir CMICB Clin069 – Gender incongruence ser	•			
9.3	Non-NHS Drug and Alcohol Rehabilitation (non- NHS commissioned services)	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin072 – Private Drug and Alcohol Rehabilitation				
9.4	Private Mental Health (MH) Care - Non-NHS Commissioned Services: including Psychotherapy, adult eating disorders, general in-patient care,post- traumatic stress adolescent mental health	This policy has been superseded by NHS Wir CMICB Clin073 – Mental health disorders, s				

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments		
10. N	0. Neurology					
10.1	Bobath Therapy	This policy has been superseded by NHS Wir CMICB Clin063 – Bobath Therapy	rral & Merseyside ICB Policy:			
10.2	Trophic Electrical Stimulation for Facial/Bells Palsy	This policy has been superseded by NHS Wir CMICB Clin062 – Idiopathic Facial Paralysis (
10.3	Functional Electrical Stimulation (FES)	This policy has been superseded by NHS Wir CMICB Clin064 – Foot Drop, Functional Elect				
11. C	phthalmology					
11.1	Upper Lid Blepharoplasty - Surgery on the Upper Eyelid	Only commissioned in the following circumstances: Eyelid function interferes with visual field.	Eyelid Surgery The British Association of Aesthetic Plastic Surgeons 2011. Modernisation Agency's Action on Plastic Surgery 2005. Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base London Health Observatory 2010.	Excess skin in the upper eyelids can accumulate due to the ageing and is thus normal. Hooded lids causing significant functional impaired vision confirmed by an appropriate specialist can warrant surgical treatment.		
				Impairment to visual field to be documented.		
11.2	Lower Lid Blepharoplasty - Surgery on the	Only commissioned in any of the following circumstances:	Eyelid Surgery The British Association of Aesthetic Plastic Surgeons 2011.	Excessive skin in the lower lid may cause "eye bags" but does not affect function		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
11.3	Surgical Treatments for Xanthelasma Palpebrum (fatty	 Correction of ectropion or entropion which threatens the health of the affected eye. Removal of lesions of eyelid skin or lid margin. Rehabilitative surgery for patients with thyroid eye disease. This policy has been superseded by NHS Wire CMICB Clin005 – Benign skin lesions	Local PCT consensus – review conducted 2007. Modernisation Agency's Action on Plastic Surgery 2005. Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010. Tral & Merseyside ICB Policy:	of the eyelid or vision and therefore does not need correction.
11.4	deposits on the eyelids) Surgery or Laser Treatment for Short Sightedness (myopia) or Long Sightedness (hypermetropia)	This policy has been superseded by NHS Wir CMICB Clin034 - Myopia, Hyperopia and Ast		
11.5	Cataract Surgery	This policy has been superseded by NHS Wir CMICB Clin097 – Cataract Surgery	rral & Merseyside ICB Policy:	
11.6	Coloured (irlens) Filters for Treatment of Dyslexia	This policy has been superseded by NHS Wir CMICB Clin017 - Dyslexia Treatment using C		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments	
11.7	Intra Ocular Telescope for Advanced Age- Related Macular Degeneration		This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin003 - Age-Related Macular Degeneration (AMD), implantable miniature telescope (IMT)		
11.8	Surgical Removal of Chalazion or Meibomian Cysts		This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin011 - Chalazia (meibomian cysts), removal		
11.9	Surgical treatment for Proptosis/ Dysthyroid eye disease	Only commissioned if:	http://patient.info/doctor/thyroid-eye-disease-pro		
11.10	Photodynamic Therapy for ARMD	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin079 – Age Related Macular Degeneration (AMD)- Photodynamic Therapy			
11.11	Multifocal (non- accommodative) intraocular lenses	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin003 - Age-Related Macular Degeneration (AMD), implantable miniature telescope (IMT)			
12. C	ral Surgery				
12.1	Surgical Replacement of the Temporo- Mandibular Joint	This policy has been superseded by NHS Wir CMICB Clin093 – Temporomandibular joint,			
	Temporo- Mandibular Joint Dysfunction Syndrome & Joint Replacement				

WIRCCG_2024 – Commissioning Policy Version 5, 18 March 2024

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments		
13. P	3. Paediatrics					
13.1	Cranial Banding for Positional Plagiocephaly	This policy has been superseded by NHS W CMICB Clin039 - Positional Plagiocephaly/b	•			
14. P	lastic & Cosmetic					
14.1	Reduction Mammoplasty - Female Breast Reduction	This policy has been superseded by NHS W CMICB Clin007 – Breast Reduction	irral & Merseyside ICB Policy:			
14.2	Augmentation Mammoplasty - Breast Enlargement	This procedure is not routinely commissioned. The following exceptions apply: In all cases: • The BMI is <25 and stable for at least twelve months. AND • Congenital absence i.e. no obvious breast tissue. In special circumstances reconstructive surgery may be appropriate for tubular breast abnormality. Patients requiring reconstructive surgery post cancer treatment are excluded from this policy. All non-surgical options must have been explored e.g. padded bra. Non-core procedure Interim Gender	Dixon, J, et al, 1994, ABC of breast diseases: congenital problems and aberrations of normal breast development and involution, Br Med J, 309, 24 September, 797-800 . Freitas, R, et al, 2007, Poland's Syndrome: different clinical presentations and surgical reconstructions in 18 cases, Aesthet Plast Surg, 31, 140-46. Heimberg, D, et al, 1996, The tuberous breast deformity: classification and treatment, Br J Plast Surg, 49, 339-45. Pacifico, M, et al, 2007, The tuberous breast revisited, J Plast Reconstruct Aesthet Surg, 60, 455-64. North Derbyshire, South Derbyshire and Bassetlaw Commissioning Consortium, 2007, Norcom commissioning policy —	Patients should be made aware that: 1 in 5 implants need replacing within 10 years regardless of make. Prior to implant insertion all patients explicitly be made aware of the possibilities of complications, implant life span, the need for possible removal of the implant at a future date and that future policy may differ from current policy. Patients should be made aware that implant removal in the future might not be automatically followed by replacement of the implant.		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		Dysphoria Protocol & Service Guidelines 2013/14.	specialist plastic surgery procedures", 5-7. Sadove, C, et al, 2005, Congenital and acquired pediatric breast anomalies: a review of 20 years experience, Plast Reconstruct Surg, April, 115(4), 1039-1050.	Not all patients demonstrate improvement in psychosocial outcome measures following breast augmentation.
			Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010.	
			Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service	
			Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013).	
			Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	
14.3	Removal and/or Replacement of Silicone Implants - Revision of Breast	Revisional surgery will ONLY be considered if the NHS commissioned the original surgery and complications arise which necessitates surgical intervention.	Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010.	1 in 5 implants need replacing within 10 years regardless of make.

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	Augmentation	If revisional surgery is being carried out for implant failure, the decision to replace the implant(s) rather than simply remove them will be based upon the clinical need for replacement and whether the patient meets the policy for augmentation at the time of revision. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core" surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG.	Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service Poly Implant Prothèse (PIP) breast implants: final report of the Expert Group Department of Health (June 2012). Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013). Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	Prior to implant insertion all patients explicitly be made aware of the possibilities of complications, implant life span, the need for possible removal of the implant at a future date and that future policy may differ from current policy. Patients should be made aware that implant removal in the future might not be automatically followed by replacement of the implant.
14.4	Mastopexy - Breast Lift	This policy has been superseded by NHS Win CMICB Clin030 – Mastopexy (breast lift)		
14.5	Surgical Correction of Nipple Inversion	This policy has been superseded by NHS Win CMICB Clin035 – Nipple inversion, surgical of		
14.6	Male Breast Reduction Surgery for Gynaecomastia	Not routinely commissioned. The following exception will apply:	Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010.	Ensure breast cancer has been excluded as a possible cause especially if there is a family history of

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		gynaecomastia caused by cancer treatment Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core" surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG.	Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service Dickson, G. (2012). Gynecomastia. American Family Physician, 85(7), 716–722. Retrieved from: http://www.aafp.org/afp/2012/0401/p716.pdf Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013). Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	breast cancer.
14.7	Hair Removal Treatments including Depilation Laser Treatment or Electrolysis – for Hirsutism	Routinely commissioned in the case of those undergoing treatment for pilonidal sinuses to reduce recurrence. In other circumstances not routinely commissioned. Will be considered via Individual Funding Request if all of the following clinical circumstances are met; • Abnormally located hair-bearing skin following reconstructive surgery located on face and neck.	Epidemiology, diagnosis and management of hirsutism: a consensus statement by the Androgen Excess and Polycystic Ovary Syndrome Society. Escobar et al. Human Reproduction Update, 03-04 2012, vol./is. 18/2(146-70). cks.nice.org.uk/hirsutism#!scenario - NICE: Clinical Knowledge Summaries 2010. Laser and photoepilation for unwanted hair	The method of depilation (hair removal) considered will be the most appropriate form usually diathermy, electrolysis performed by a registered electrologist, or laser centre.

Treatment / Procedure	Eligibility Criteria	Evidence	Comments
a a a s e e e e e e e e e e e e e e e e	There is an existing endocrine medical condition and severe facial hirsutism. 1. Ferryman Gallwey (<i>A method of evaluating and quantifying hirsutism in women</i>) Score 3 or more per area to be treated. 2. Medical treatments have been tried for at least one year and failed. 3. Patients with a BMI of>30 should be in a weight reduction programme and should have lost at least 5% body weight. All cases will be subject to individual approval by the IFR Team and must be accompanied by an opinion from a secondary care consultant (i.e. endocrinologist). Photographs will also be required to allow the CCG's to visibly asses the severity equitably. Funded for 6 treatments only at an NHS commissioned premises. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core"	Management of hirsutism – Koulouri et al BMJ 2009; 338:b847. Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013). Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG.		
14.8	Surgical Treatment for Pigeon Chest	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: <u>CMICB_Clin038 – Pectus Deformity, surgical treatment</u>		
14.9	Surgical Revision of Scars	Funding of treatment will be considered only for scars which interfere with function following burns, trauma, treatments for keloid, or post-surgical scarring. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core" surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG.	Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013). Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	
14.10	Laser Tattoo Removal	 Only commissioned in any of the following circumstances: Tattoo is result of trauma inflicted against the patient's will. The patient was a child and not responsible for his/her actions at the time of tattooing. Inflicted under duress. During adolescence or disturbed periods (only in very exceptional circumstances) 	Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010. Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		 where tattoo causes marked limitations of psycho-social function). An individual funding request will be required. 	Modernisation Agency's Action on Plastic Surgery 2005.	
14.11	Apronectomy or Abdominoplasty (Tummy Tuck)	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin099 – Abdominoplasty or Apronectomy (tummy tuck)		
14.12	Other Skin Excisions/ Body Contouring Surgery e.g. Buttock Lift, Thigh Lift, Arm Lift (Brachioplasty)	This policy has been superseded by NHS Wir CMICB Clin006 – Body Contouring and other		and arm lift (brachioplasty)
14.13	Treatments to Correct Hair Loss for Alopecia	Only commissioned in either of the following circumstances: Result of previous surgery. Result of trauma, including burns. Hair Intralace System is not commissioned. Dermatography is not commissioned. NHS wigs will be available according to NHS policy. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core"	British Association of Dermatologists' guidelines for the management of alopecia areata 2012 Interventions for alopecia areata — Cochrane Library 2008. http://www.bad.org.uk/library-media%5Cdocuments%5CAlopecia areat a guidelines 2012.pdf Only one study which compared two topical corticosteroids showed significant short-term benefits. No studies showed long-term beneficial hair growth. None of the included studies asked participants to report their opinion of hair growth or whether their quality of life had improved with the treatment.	

Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG	No evidence of effective treatments for alopecia – Cochrane Pearls 2008. Alopecia areata – NICE Clinical Knowledge Summaries 2008. Health Commission Wales. 2008 Commissioning Criteria – Plastic Surgery. Procedures of Low Clinical Priority/ Procedures not usually available on the National Health Service Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010 (further evidence provided within this document by Islington PCT to support funding). Modernisation Agency's Action on Plastic Surgery 2005. Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013). Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
14.14	Hair Transplantation	Commissioned only in exceptional circumstance, e.g. reconstruction of the eyebrow following cancer or trauma. Dermatography may be an acceptable alternative in eyebrow reconstruction. Non-core procedure Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. Where the provision of "non-core" surgeries is appropriate, the GIC should apply for treatment funding through the CCG; the GIC should endeavour to work in partnership with the CCG.	A trial on subcutaneous pedicle island flap for eyebrow reconstruction – Mahmood & Mehri. Burns, 2010, Vol. 36(5), p692-697. Procedures of Limited Clinical Effectiveness Phase 1 - Consolidation and repository of the existing evidence-base - London Health Observatory 2010 further evidence provided within this document by Islington PCT to support funding. Modernisation Agency's Action on Plastic Surgery 2005. Interim Gender Dysphoria Protocol & Service Guidelines 2013/14. NHS England interim protocol NHS England (2013). Pages 13 & 14 describe non-core NHS England & CCG commissioning responsibilities.	
14.15	Treatments to Correct Male Pattern Baldness	This is not routinely commissioned.	Modernisation Agency's Action on Plastic Surgery 2005.	
14.16	Labiaplasty, Vaginoplasty and Hymenorrhaphy	This policy has been superseded by NHS Win CMICB Clin077 – Labiaplasty, vaginoplasty		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments			
14.17	Liposuction	This policy has been superseded by NHS Wirral & Merseyside ICB Policy:					
		CMICB Clin0100 – Liposuction	CMICB Clin0100 – Liposuction				
14.18	Rhytidectomy - Face or Brow Lift	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:				
	or Brow Lift	CMICB Clin042 - Rhytidectomy	CMICB Clin042 – Rhytidectomy				
14.19	All procedures	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:				
	undertaken on cosmetic grounds	CMICB Clin013 – Cosmetic Procedures					
	espiratory						
15.1							
	Sodium Tetradecyl Sulfate (STS) Injection or 'snoreplasty'						
	Uvulopalatoplasty and						
	Uvulopalatopharyng opl asy						
15.2	Investigations and	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:				
	treatment for Sleep Apnoea	CMICB Clin074 – Sleep Apnoea or Narcole	psy referral and management				

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
15.3	Sleep studies/ Hypersomnia	This policy has been superseded by NHS Wir CMICB Clin074 – Sleep Apnoea or Narcole		
16. 1	Low back pain and sciatica in over 16's Diagnostic, Interventions and Treatments for acute and chronic low back pain. Excluding spinal pathology, radiculopathy and children. Pharmacological Intervention for lower back pain Pharmacological intervention for sciatica (neuropathic pain in adults)	Diagnostic imaging should not be routinely offered unless: • in a specialist setting where the results are likely to change clinical management OR • Diagnostic imaging is required prior to referral for surgical intervention Management should be in line with NICE Guidance and should consist of advice and information to enable self-management. Patients should be encouraged to continue with normal activities. Structured exercise programmes (including group exercise), psychological therapies and manual therapy should be considered. Manual therapy should only be offered as part of a treatment package, including exercise with or without psychological therapy. Spinal injections of local anaesthetic and	https://www.nice.org.uk/guidance/NG59	Amendments made based on version developed by Lancashire & Midlands CSU in collaboration with the Walton Centre

Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	steroid should not be offered for patients with non-specific low back pain.		
	For people with non-specific low back pain the following injections should not be offered:		
	Facet joint injections Therapeutic medial branch blocks Intradiscal therapy Prolotherapy Trigger point injections with any agent, including botulinum toxin Epidural steroid injections for chronic low back pain or for neurogenic claudication in patients with central spinal canal stenosis		
	Any other spinal injections not specifically covered above Radiofrequency denervation can be offered according to NICE guideline (NG59) if all non-surgical and alternative treatments have been tried and there is moderate to severe chronic pain that has improved in response to diagnostic medical branch block.		
	Epidurals (local anaesthetic and steroid) should be considered in patients who have acute and severe lumbar radiculopathy at time of referral. Alternative and less invasive options have been shown to work e.g. exercise programmes, behavioural therapy,		

Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	and attending a specialised pain clinic. Alternative options are suggested in line with the National Back Pain Pathway. Consider oral non-steroidal anti-inflammatory drugs (NSAIDS) at lowest effective dose for shortest possible time. If NSAIDS are contraindicated/ not tolerated or ineffective, consider weak opiods. Do not offer: Paracetamol alone Opioids for acute low back pain (unless NSAIDs are contraindicated) Opioids for chronic low back pain Selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine, reuptake inhibitors or	Evidence	Comments
	 tricyclic antidepressants Anticonvulsants Offer a choice of amitriptyline, duloxetine, gabapentin or pregabalin as initial treatment for neuropathic pain (except trigeminal neuralgia). If the initial treatment is not effective or is not tolerated, offer one of the remaining 3 drugs, and consider switching again if the second and third drugs tried are also not effective or not tolerated. 		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		Consider tramadol only if acute rescue therapy is needed (see NICE Guidance CG173 for long term use). Consider capsaicin cream[4] for people with localised neuropathic pain who wish to avoid, or who cannot tolerate, oral treatments. Do not start the following to treat neuropathic pain in non-specialist settings, unless advised by a specialist to do so: cannabis sativa extract capsaicin patch lacosamide lamotrigine levetiracetam morphine oxcarbazepine topiramate tramadol (this is referring to longterm use) venlafaxine		
16.2	Radiofrequency Facet Joint Denervation	This policy has been superseded by NHS Wir CMICB Clin089 – Chronic Low Back Pain, I	•	,
16.3	Fusion	This policy has been superseded by NHS Wir CMICB Clin087 – Spinal fusion surgery for		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
16.4	Epidural Injection	This policy has been superseded by NHS Wir CMICB Clin060 – Spinal Injections for Low Ba	•	
16.5	Spinal Decompression	Consider spinal decompression for people with sciatica when non-surgical treatment has not improved pain or function and their radiological findings are consistent with sciatic symptoms.		
16.6	Endoscopic Laser Foraminoplasty	This policy has been superseded by NHS Wir CMICB Clin018 – Endoscopic Laser Foramin		
16.7	Peripheral Nerve- field Stimulation (PNFS) for Chronic Low Back Pain	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin012 – Chronic Low Back Pain, Peripheral Nerve Field Stimulation		
16.8	Endoscopic Lumbar Decompression	This procedure is NOT routinely commissioned.	IPG300: Percutaneous endoscopic laser lumbar discectomy NICE, 2009	
16.9	Percutaneous Disc Decompression using Coblation for Lower Back Pain	This procedure is NOT routinely commissioned.	IPG 173: Percutaneous disc decompression using coblation for lower back pain. NICE 2006	
16.10	Non-Rigid Stabilisation Techniques	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin090 – Non-rigid stabilisation techniques for degenerative disease of the lumbar spine		
16.11	Lateral (including extreme, extra and direct lateral) Interbody Fusion in	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin087 – Spinal fusion surgery for non-specific, mechanical back pain		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	the Lumbar Spine			
16.12	Percutaneous Intradiscal Laser Ablation in the Lumbar Spine	This procedure is NOT routinely commissioned.	IPG 357: Percutaneous intradiscal laser ablation in the lumbar spine NICE 2010.	
16.13	Transaxial Interbody Lumbosacral Fusion	This policy has been superseded by NHS Wir CMICB Clin087 – Spinal fusion surgery for		
16.14	Therapeutic Endoscopic Division of Epidural Adhesions	This policy has been superseded by NHS Wir CMICB_Clin019 – Epidural Adhesions, Thera		
16.15	Automated Percutaneous Mechanical Lumbar Discectomy	This procedure is NOT routinely commissioned.	IPG 141: <u>Automated percutaneous</u> mechanical lumbar discectomy. Nov 2005.	
16.16	Prosthetic Intervertebral Disc Replacement in the Lumbar Spine	This policy has been superseded by NHS Wir CMICB Clin029 – Low back pain, disc replacen	•	
16.17	Bone Morphogenetic Proteins - Dibotermin Alfa; Eptotermin Alpha	Dibotermin alfa is commissioned in the following situation: The treatment of acute tibia fractures in adults, as an adjunct to standard care using open fracture reduction and intramedullary unreamed nail fixation.	Clinical effectiveness and cost- effectiveness of bone morphogenetic proteins in the non-healing of fractures and spinal fusion: a systematic review Health Technology Assessment NHS R&D HTA Programme, 2007.	
		Eptotermin alfa is commissioned in line with	Clinical effectiveness and cost-effect [Health Technol Assess. 2007] - PubMed -	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		its licensed indication: Treatment of non-union of tibia of at least 9 month duration, secondary to trauma, in skeletally mature patients, in cases where previous treatment with autograft has failed or use of autograft is unfeasible.	Annals of Internal Medicine Safety and Effectiveness of Recombinant Human Bone Morphogenetic Protein-2 for Spinal Fusion: A Meta-analysis of Individual-Participant Data June 2013 BMPs: Options, indications, and effectiveness – Journal of Orthopaedic Trauma. 2010 Mar;24 Suppl 1:S9-16.	
16.18	Surgery for Trigger Finger	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin048 – Trigger Finger release in adults		
16.19	Hyaluronic Acid and Derivatives Injections for Peripheral Joint Pain	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin036 – Osteoarthritic induced changes in peripheral joints (knee, hips, ankle & thumb), intra-articular hyaluronan (hyaluronic acid)		
16.20	Secondary Care Administered Steroid Joint Injections	This policy has been superseded by NHS Win CMICB Clin037 – Osteoarthritis-induced join		articular corticosteroids
16.21	Dupuytren's Disease Palmar Fasciectomy/Needle Faciotomy Radiotherapy	This policy has been superseded by NHS Win CMICB_Clin016 - Dupuytren's Contracture		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments	
	Collagenase Injections for Dupuytren's Disease				
	Dupuytren's Disease Surgical treatment				
	Dupuytrens				
	Contracture – conservative treatment				
16.22	Hip and Knee Replacement	This policy has been superseded by NHS Wirral & Merseyside ICB Policy:			
	Surgery & Hip Resurfacing	CMICB Clin084 – Hip and knee replacement surgery			
16.24	Diagnostic Arthroscopy for	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:		
	Arthritis of the Knee	CMICB Clin004 – Arthroscopic Surgery of the	e Knee for Meniscal Tears		
16.25	Arthroscopic Lavage and	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:		
	Debridement for	CMICB Clin028 – Knee Osteoarthritis, Arthroscopic Lavage and Debridement			
	Osteoarthritis of the Knee				
16.26	Patient Specific Total Knee	This policy has been superseded by NHS Wir	ral & Merseyside ICB Policy:		
	Replacement	CMICB Clin047 – Total Knee Arthroplasty, pa	atient specific instrumentation/implants		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments	
16.27	Surgical Treatment for Carpal Tunnel Syndrome	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin010 – Carpal Tunnel interventions and surgery			
16.38	Nerve Conduction Studies for Carpal Tunnel Syndrome	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin112 – Carpal Tunnel Syndrome, Nerve Conduction Testing			
16.29	Surgical Removal of Mucoid Cysts at Distal Inter Phalangeal Joint (DIP)	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB_Clin033 – Mucoid Cysts of the Fingers at the Distal Interphalangeal (DIP) Joint, surgical removal			
16.30	Surgical Removal of Ganglions	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: <u>CMICB_Clin022 – Ganglia, surgical removal and general management</u>			
16.31	Hip Arthroscopy for Femoro–Acetabular Impingement	CCGs routinely commission hip arthroscopy (from surgeons with specialist expertise in this type of surgery) in line with the requirements stipulated by NICE IPG 408, and only for patients who fulfil ALL of the following criteria: A definite diagnosis of hip impingement syndrome/femoro-acetabular impingement (FAI) has been made by appropriate investigations, X-rays, MRI and CT scans.	IPG408 Arthroscopic femoro-acetabular surgery for hip impingement syndrome: guidance – NICE, 2011. http://www.hullccg.nhs.uk/uploads/policy/file/22/hip-arthroscopy-hull-ccg.pdf NHS Hull Clinical Commissioning Group 2012. Vijay D Shetty, Richard N Villar. Hip arthroscopy: current concepts and review of literature. British Journal of Sports Medicine, 2007;41:64–68.	Current evidence on the efficacy of arthroscopic femoro–acetabular surgery for hip impingement syndrome is adequate in terms of symptom relief in the short and medium term. With regard to safety, there are well-recognised complications. Therefore	
		An orthopaedic surgeon who specialises in young adult hip surgery has made the diagnosis in collaboration with a specialist	Macfarlane RJ, Haddad FS The diagnosis and management of femoro-acetabular impingement. Annals of the Royal College of Surgeons of England, July 2010, vol/iss	this procedure may be used provided that normal arrangements are in place for clinical governance,	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		musculoskeletal radiologist. The patient has had severe FAI symptoms (restriction of movement, pain and 'clicking') or significantly compromised functioning for at least 6 months. The symptoms have not responded to all available conservative treatment options including activity modification, drug therapy (NSAIDs) and specialist physiotherapy.	92/5(363-7). Ng V Y et al Efficacy of Surgery for Femoro-acetabular Impingement: A Systematic Review. American Journal of Sports Medicine, November 2010,38 2337-2345. Commissioning Guide: Painful osteoarthritis of the hip Royal College of Surgeons (2013). IPG408 Arthroscopic femoro-acetabular surgery for hip impingement syndrome: guidance NICE, 2011	consent and audit with local review of outcomes.
16.32	Surgical Removal of Bunions/Surgery for Lesser Toe Deformity	This policy has been superseded by NHS Wir CMICB_Clin008 – Bunions, surgical removal	ral & Merseyside ICB Policy:	
16.33	Surgical Treatment of Morton's Neuroma	This policy has been superseded by NHS Wir CMICB Clin032 – Morton's Neuroma, surgical		
16.34	Surgical Treatment of Plantar Fasciitis	Surgical Treatment is not routinely commissioned unless the following pathway has been followed: 1. Patient has documented evidence that they are not responding to conservative treatments 2. Patient is experiencing significant pain or it is having a serious impact on their	Heel painplantar fasciitis: clinical practice guidelines linked to the international classification of function, disability, and health from the orthopaedic section of the American Physical Therapy Association - Journal of Orthopaedic & Sports Physical Therapy. 2008:38(4):A1-A18.	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
16.35	Treatment of Tendinopathies Extracorporeal Shock Wave Therapy Autologous Blood or Platelet Injection	daily life and has completed the following. 3. Three months of conservative therapy such as footwear modification, stretching exercises, ice packs, weight loss. 4. Been referred to a podiatrist or physiotherapist. 5. Not responded to corticosteroid injections. This policy has been superseded by NHS Wir CMICB Clin001 - Achilles Tendinopathy, Ref shockwave therapy, autologous blood or plate	ractory Tennis Elbow and Plantar Fasciitis: tre	eatment with extracorporeal
16.36	Injection Injections for Tendonitis (Jumpers Knee)	Injections for Tendonitis (Jumpers Knee) are not routinely commissioned.	http://www.nhs.uk/Conditions/Tendonitis/Pages/Treatment.aspx	
16.37	Shoulder Arthroscopy (including arthroscopic shoulder decompression for subacromial shoulder pain)	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin092 – Subacromial shoulder pain, arthroscopic shoulder decompression surgery		
16.38	Hip Injections	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: <u>CMICB Clin059 – Hip pain, intra-articular injections of corticosteroids</u>		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments		
17. L	7. Urology					
17.1 Circumcision This policy has been superseded by NHS Wirral & Merseyside ICB Policy:						
	CMICB Clin104 – Penile circumcision in children and young people under 16 years					
17.2	Penile Implant: A Surgical Procedure	This policy has been superseded by NHS Wir	rral & Merseyside ICB Policy:			
	to Implant a Device	CMICB_Clin020 - Erectile dysfunction, penile	e prosthesis surgery			
17.3	Male sterilisation under General Anaesthetic	Not routinely commissioned.				
17.4	Reversal of Male Sterilisation	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: ICB Policy CMICB_Clin040 – Reversal of Male Sterilisation				
17.5	ESWT (extracorporeal	This policy has been superseded by NHS Wirral & Merseyside ICB Policy:				
	shockwave therapy)	CMICB Clin111 – Chronic Pelvic Pain Syndrome in Men, Hyperthermia, Extracorporeal Shockwave Therapy				
	for Prostadynia or	and Sacral Neuromodulation				
	Pelvic Floor Syndrome	and odoral Hodromoddiadon				
17.6	Hyperthermia Treatment for	This policy has been superseded by NHS Wirral & Merseyside ICB Policy:				
	Prostadynia or	CMICB Clin111 - Chronic Pelvic Pain Synd	lrome in Men, Hyperthermia, Extracorpore	eal Shockwave Therapy		
	Pelvic Floor Syndrome	and Sacral Neuromodulation				
17.7	Surgery for Prostatism	Only commissioned where there are sound clinical reasons and after failure of conservative treatments and in any of the following circumstances: International prostate symptom score >7;	CG97: Lower urinary tract symptoms: The management of lower urinary tract symptoms in men NICE 2010.	No references to treatment thresholds found.		

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
17.8	Surgical treatment for Hydroceles – adults and children	 dysuria; Post voided residual volume >150ml; Recurrent proven Urinary Tract Infections (UTI); Deranged renal function; Prostate-specific antigen (PSA) > age adjusted normal values. Only commissioned if: In the case of communicating hydrocele: patient is aged over 18 months of age In the case of non-communicating hydrocele, the patient is experiencing: discomfort and/or disfigurement resulting in functional impairment preventing individual fulfilling work/study/carer or domestic duties (adult) or discomfort and/or disfigurement resulting in inability to participate in normal social and educational activity (adolescent) 	LUTS in men, age-related (prostatism) NICE Clinical Knowledge Summaries (2010). http://www.rcseng.ac.uk/healthcare- bodies/docs/published-guides/luts Royal College of Surgeons (2013). http://patient.info/health/hydrocele-in- adults	
17.9	Surgical removal of benign epididymal cysts	Only commissioned if ALL the following criteria are met: it is large enough to cause change in shape and size of scrotum cyst is putting pressure on other structures in the testes cyst is causing prolonged or significant pain	http://patient.info/health/epididymal-cyst	

	Treatment / Procedure	Eligibility Criteria	Evidence	Comments		
18. V	18. Vascular					
18.1 Surgery for Extreme Sweating		This policy has been superseded by NHS Wil				
	Hyperhydrosis – all areas	CMICB Clin027 – Hyperhidrosis (excessive sweating), Surgical Management				
	Surgical Resection Endoscopic Thoracic Sympathectomy					
18.2	Chelation Therapy for Vascular Occlusions	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin015 – Disodium Ethylenediaminetetraacetic Acid (EDTA) in prevention of Cardiovascular Events in patients with a previous Myocardial Infarction				
18.3	Varicose Veins Interventional Treatments e.g. endothermal ablation, foam sclerotherapy and surgery	This policy has been superseded by NHS Wirral & Merseyside ICB Policy: CMICB Clin049 – Varicose Veins				
19. C						
19.1	Botulinum Toxin A & B Used in several types of procedures e.g. to treat muscle disorders, excessive sweating hyperhidrosis) and migraine.	The use of botulinum toxin type A is commissioned in the following indications: • Anal fissures only following a minimum of two months with standard treatment (lifestyle and topical pharmaceutical products) for chronic anal fissures that have not resulted in fissure healing; and	NICE TA260 June 2012 – Migraine (chronic) botulinum toxin type A http://guidance.nice.org.uk/TA260 Idiopathic detrusor instability - only commissioned in accordance with NICE CG171 Sept 2013 - Urinary incontinence in women			

WIRCCG_2024 – Commissioning Policy Version 5, 18 March 2024

Treatment / Procedure	Eligibility Criteria	Evidence	Comments
Procedure	only a maximum of 2 courses of injections. Blepharospasm and hemifacial spasm. Probable contracture of joint in multiple sclerosis, in conjunction with prolonged stretching modalities (i.e. in line with NICE Clinical Guideline 8). http://guidance.nice.org.uk/CG8 Focal dystonia, where other measures are inappropriate or ineffective. Focal spasticity in patients with upper motor neurone syndrome, caused by cerebral palsy, stroke, acquired brain injury, multiple sclerosis, spinal cord injuries and neurodegenerative disease, where other measures are inappropriate or ineffective. Idiopathic cervical dystonia (spasmodic torticollis). Prophylaxis of headaches in adults with chronic migraine (defined as headaches on at least 15 days per month of which at least 8 days are with migraine) that has not responded to at least three prior pharmacological prophylaxis therapies, and whose condition is appropriately managed for medication overuse (i.e. in line with NICE Technology Appraisal 260). http://guidance.nice.org.uk/TA260 Refractory detrusitor overactivity, only line with NICE Clinical Guideline 171 (women) http://guidance.nice.org.uk/CG171 and	http://guidance.nice.org.uk/CG171 Diagnosis and management of hyperhidrosis British Medical Journal.	

Treatment / Procedure	Eligibility Criteria	Evidence	Comments
	Clinical Guideline 97 (men) http://guidance.nice.org.uk/CG97 where conservative therapy and conventional drug treatment has failed to control symptoms. • Sialorrhoea (excessive salivary drooling), when all other treatments have failed. Botulinum toxin type A is not routinely commissioned in the following indications: • Canthal lines (crow's feet) and glabellar (frown) lines. • Hyperhidrosis. • Any other indication that is not listed above		
	The use of Botulinum Type B is not routinely commissioned.		
	Where the use of botulinum toxin is used to treat an indication outside of the manufacturer's marketing authorisation, clinicians and patients should be aware of the particular governance requirements, including consent (which must be documented) for using drugs outside of their licensed indications.		
	For patients with conditions which are not routinely commissioned, as indicated above, requests will continue to be considered by Cheshire & Wirral Clinical Commissioning		


	Treatment / Procedure	Eligibility Criteria	Evidence	Comments
		Groups processes for individual funding requests, if there is evidence that the patient is considered to have clinically exceptional circumstances to any other patient experiencing the same condition within Cheshire & Wirral. Requests to commission the use of botulinum toxin as an option to treat other indications, where a known cohort of patients can be identified, should be processed in accordance with the relevant CCG's defined processes. If a subsequent CCG approved policy supersedes the information above, this section will be reviewed and updated.		
19.2	Correction of privately funded treatment	Correction of privately funded treatment is not routinely commissioned unless in an emergency.		

J. Appendix 1 – Cataract Referral Guide

This policy has been superseded by NHS Wirral & Merseyside ICB Policy:

CMICB_Clin097 - Cataract Surgery

L. Appendix 2 – IEFR Process



M. Appendix 3 – IFER Panel Contact Details

Telephone: 01244 650 305

CCG	Email Address
West Wirral CCG	WestWirralccg.IFR@nhs.net
Eastern Wirral CCG	Eastern Wirralccg.IFR@nhs.net
South Wirral CCG	southWirralccg.ifr@nhs.net
Vale Royal CCG	valeroyalccg.ifr@nhs.net

N. Appendix 4 – Fusion Surgery – Clinical exceptions permitted

This policy has been superseded by NHS Wirral & Merseyside ICB Policy:

CMICB Clin087 - Spinal fusion surgery for non-specific, mechanical back pain